

GOAT ANTI-BDH2 / DHRS6 (AA 60 TO 71) ANTIBODY

SKU: EB08290



250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym / Alias Names	oxidoreductase UCPA dehydrogenase/reductase (SDR family) member 6 UNQ6308 UCPA-OR PRO20933 FLJ13261 EFA6R 3-hydroxybutyrate dehydrogenase, type 2 DHRS6 BDH2
Accession ID	NP_064524.3
Blocking Peptide	EBP08290
Immunogen	Peptide with sequence C-TKKKQIDQFANE, from the internal region of the protein sequence according to NP_064524.3.
Peptide Sequence	C-TKKKQIDQFANE
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Rat
Reactive Species	Human
Human Gene ID	56898
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
IHC Results	Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.
ELISA Detection Limit	Antibody detection limit dilution 1:64000.
Western Blot	Approx 26kDa band observed in Human Kidney lysates (calculated MW of 26.7kDa according to NP_064524.3). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.
Application Type	Pep-ELISA, WB, IHC

SELECTED REFERENCES

[{"pmid": 21492153, "intro": "**This antibody has been successfully used in Western Blot on Human:**", "title": "Analysis of proteomic changes induced upon cellular differentiation of the"}]

human intestinal cell line Caco-2.", "author": "Buhrke T, Lengler I, Lampen A.", "journal": "Dev Growth Differ. 2011 Apr;53(3):411-26"}]

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

