

## GOAT ANTI-BLK ANTIBODY

**SKU:** EB10736

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

---

## SPECIFICATIONS

<b>Formulation</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
<b>Unit Size</b>	100 µg
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Instructions</b>	
<b>Synonym /</b>	
<b>Alias</b>	MODY11  MGC10442  B lymphoid tyrosine kinase BLK
<b>Names</b>	
<b>Accession ID</b>	NP_001706.2
<b>Blocking Peptide</b>	EBP10736
<b>Immunogen</b>	Peptide with sequence QWSPLKVSAQDKD-C, from the N Terminus (near) of the protein sequence according to NP_001706.2.
<b>Peptide Sequence</b>	QWSPLKVSAQDKD-C
<b>Purification Method</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Shipping Instructions</b>	Refrigerated
<b>Predicted Species</b>	Human
<b>Reactive Species</b>	Human
<b>Human Gene ID</b>	640
<b>Product Grade</b>	<a href="https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png">https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png</a>
<b>IHC Results</b>	Paraffin embedded Human Spleen. Recommended concentration: 3.75µg/ml.
<b>ELISA</b>	
<b>Detection Limit</b>	Antibody detection limit dilution 1:128000.
<b>Western Blot</b>	Approx 50kDa band observed in Human Thymus, Spleen and Lymph Node lysates (calculated MW of 57.7kDa according to NP_001706.2). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.
<b>Application Type</b>	Pep-ELISA, WB, IHC

## DOCUMENTS

- [Data Sheet](#)


## GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa



50kDa

37kDa

25kDa

20kDa

15kDa