

# GOAT ANTI-BLU / ZMYND10 ANTIBODY

**SKU:** EB05650



---

## SPECIFICATIONS

<b>Unit Size</b>	100 µg
<b>Synonym / Alias Names</b>	zinc finger, MYND domain-containing 10 zinc finger, MYND-type containing 10 FLU BLU BLu protein zinc finger, MYND domain containing 10 ZMYND10
<b>Accession ID</b>	NP_056980.2
<b>Blocking Peptide</b>	EBP05650
<b>Immunogen</b>	Peptide with sequence GKTCVLAAQGDRAK, from the C Terminus of the protein sequence according to NP_056980.2.
<b>Peptide Sequence</b>	GKTCVLAAQGDRAK
<b>Shipping Instructions</b>	Refrigerated
<b>Predicted Species</b>	Human, Mouse, Rat
<b>Human Gene ID</b>	51364
<b>Mouse Gene ID</b>	114602
<b>Product Grade</b>	<a href="https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/aspiring_medium.png">https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/aspiring_medium.png</a>
<b>ELISA Detection Limit</b>	Antibody detection limit dilution 1:16000.
<b>Western Blot</b>	Preliminary experiments gave an approx 75-80kDa band in A549 lysates at 1ug/ml, this band was successfully blocked by incubation with the immunising peptide. Please note that currently we cannot find an explanation in the literature for the band we observe given the predicted size of approx. 50kDa according to NP_056980. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?
<b>Application Type</b>	Pep-ELISA

## DOCUMENTS

- [Data Sheet](#)