


Telephone: (650) 697-3600

## **GOAT ANTI-C-FOS (AA283-295) ANTIBODY**

**SKU:** EB11742



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Storage

**Alias** 

Aliquot and store at -20°C. Minimize freezing and thawing. Instructions

Synonym / proto-oncogene c-Fos|p55|G0/G1 switch regulatory protein 7|FOS|FBJ murine osteosarcoma viral oncogene homolog|FBJ murine osteosarcoma viral (v-fos) oncogene homolog (oncogene FOS)|C-FOS|cellular oncogene c-

**Names** fos|AP-1|activator protein 1

Usage <strong>Immunofluorescence:</strong> Strong expression of the protein seen in the nucleus of HeLa cells,

and in the ER and cytoplasm of MCF7 cells. Recommended concentration:  $10\mu g/ml$ . **Summary** 

Accession

NP\_005243.1 ID

**Blocking** 

EBP11742 **Peptide** 

**Immunogen** 

Peptide with sequence C-SETARSVPDMDLS, from the internal region of the protein sequence according to NP 005243.1.

**Peptide** 

C-SETARSVPDMDLS Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.









Shipping Refrigerated Instructions

**Predicted** Human, Dog, Pig, Cow **Species** 

Reactive Human **Species** 

Human 2353 **Gene ID** 

**Product** Grade

 $https://prod-vector-labs-pimcore-assets.s 3. us-east-1. amazonaws.com/assets/products/image/elite\_medium.png$ 

IHC Results Paraffin embedded Human Tonsil. Recommended concentration: 8µg/ml.

**ELISA** 

Antibody detection limit dilution 1:128000. Detection

Limit

Type

**Application** Pep-ELISA, IF, IHC

## **GALLERY IMAGES**

