

GOAT ANTI-CD20 / MS4A1 (C TERMINUS) ANTIBODY

SKU: EB05140

▷ 05140-S4C2G2, 0.05, HLY

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa



SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym / Alias Names	MS4A2 leukocyte surface antigen Leu-16 CVID5 B-lymphocyte cell-surface antigen B1 CD20 receptor CD20 antigen MGC3969 LEU-16 CD20 BP35 S7 B1 membrane-spanning 4-domains, subfamily A, member 1 MS4A1
Accession ID	NP_068769.2; NP_690605.1
Blocking Peptide	EBP05140
Immunogen	Peptide with sequence CQDQESSPIENDSSP, from the C Terminus of the protein sequence according to NP_068769.2; NP_690605.1.
Product Comments	Please note that both reported variants (NP_068769.2 and NP_690605.1) encode the same protein.
Peptide Sequence	CQDQESSPIENDSSP
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human
Reactive Species	Human
Human Gene ID	931
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:16000.
Western Blot	Approx 35kDa band observed in Human Lymph node lysates (calculated MW of 33.1kDa according to NP_068769.2). Recommended concentration: 0.05-0.3µg/ml. Primary incubation 1 hour at room temperature.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

A 05140-S4C2G2, 0.05, HLY

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

