

GOAT ANTI-CLEC4C / BDCA2 ANTIBODY

SKU: EB09343

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
	CLEC4C C-type lectin domain family 4, member
Synonym / Alias	C BDCA2 CD303 CLECSF11 CLECSF7 DLEC HECL MGC125789 MGC125791 MGC125792 MGC125793 PRO34150 C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 11 C-type (calcium dependent, carbohydrate-recognition domain) lectin, superfamily member 7 C-type lectin domain family 4, member C, isoform 1 blood dendritic cell antigen 2 protein dendritic cell lectin b
Names	
Accession ID	NP_569708.1; NP_987099.1
Blocking Peptide	EBP09343
Immunogen	Peptide with sequence QSWTKSQKNCSV, from the Internal region of the protein sequence according to NP_569708.1; NP_987099.1.
Product Comments	This antibody is expected to recognize isoform 1 and 2 (NP_569708.1; NP_987099.1).
Peptide Sequence	QSWTKSQKNCSV
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human
Reactive Species	Human
Human Gene ID	170482
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:8000.
Western Blot	Approx 20kDa band observed in Human Spleen lysates (calculated MW of 21.5kDa according to NP_987099.1). Recommended concentration: 1-3µg/ml.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa