

# GOAT ANTI-CXCL2 / MIP-2 ANTIBODY

**SKU:** EB09352

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa

## SPECIFICATIONS

<b>Formulation</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
<b>Unit Size</b>	100 µg
<b>Storage Instructions</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Synonym / Alias Names</b>	melanoma growth stimulatory activity beta MGSA beta GRO2 oncogene SCYB2 MIP2A MIP2 MIP-2a MGSA-b GROb GRO2 CINC-2a chemokine (C-X-C motif) ligand 2 CXCL2
<b>Accession ID</b>	NP_002080.1
<b>Blocking Peptide</b>	EBP09352
<b>Immunogen</b>	Peptide with sequence C-KKIIIEKMLKNGKSN, from the C Terminus of the protein sequence according to NP_002080.1.
<b>Peptide Sequence</b>	C-KKIIIEKMLKNGKSN
<b>Purification Method</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Shipping Instructions</b>	Refrigerated
<b>Predicted Species</b>	Human
<b>Reactive Species</b>	Human
<b>Human Gene ID</b>	2920
<b>Product Grade</b>	<a href="https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png">https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png</a>
<b>ELISA Detection Limit</b>	Antibody detection limit dilution 1:32000.
<b>Western Blot</b>	Approx 15kDa band observed in in lysates of cell line Daudi (calculated MW of 11.4kDa according to NP_002080.1). Recommended concentration: 0.03-0.1µg/ml. Approx 17Da band observed in Human Duodenum lysates (calculated MW of 11.4kDa according to NP_002080.1).
<b>Application Type</b>	Pep-ELISA, WB

## DOCUMENTS

- [Data Sheet](#)

## GALLERY IMAGES

