


GOAT ANTI-CYSTATIN 8 (AA 100-112) ANTIBODY

SKU: EB09823



SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	CRES cystatin 8 cystatin 8 (cystatin-related epididymal specific) cystatin-related epididymal spermatogenic
Alias	protein cystatin-related epididymal-specific CST8
Names	
Usage	Purchase the +ve control! The transfected lysate used in QC is available for sale. Please email
Summary	sales@everestbiotech.com to request datasheet, pricing and delivery information.
Accession ID	NP_005483.1
Blocking Peptide	EBP09823
Immunogen	Peptide with sequence STNEICAIQENSK, from the internal region of the protein sequence according to NP_005483.1.
Peptide Sequence	STNEICAIQENSK
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human
Reactive Species	Human
Human Gene ID	10047
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:16000.
Western Blot	In transfected HEK293 transiently expressing Human CST8 (MYC and DYKDDDDK tagged), a band of approx. 23kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc tag antibody. Recommended concentration, 1-3µg/ml.
Application Type	Pep-ELISA, WB-Trf

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

