
Telephone: (650) 697-3600







GOAT ANTI-CYSTATIN 8 (AA 100-112) ANTIBODY



Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing. Instructions

Synonym /

Names

Alias

CRES|cystatin 8|cystatin 8 (cystatin-related epididymal specific)|cystatin-related epididymal spermatogenic

protein|cystatin-related epididymal-specific|CST8

Usage

Purchase the +ve control! The transfected lysate used in QC is available for sale. Please email

Summary

sales@everestiotech.com to request datasheet, pricing and delivery information.

Accession

NP_005483.1

Blocking

ID

EBP09823

Peptide

Peptide with sequence STNEICAIQENSK, from the internal region of the protein sequence according to

Immunogen NP 005483.1.

Peptide

Sequence

STNEICAIQENSK

Method

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

using the immunizing peptide.

Shipping

Instructions

Refrigerated

Human

10047

Predicted

Species

Reactive

Human **Species**

Human

Gene ID

Product

Grade

ELISA Antibody detection limit dilution 1:16000.

Detection Limit

Western **Blot**

In transfected HEK293 transiently expressing Human CST8 (MYC and DYKDDDDK tagged), a band of approx. 23kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png

anti-myc tag antibody. Recommended concentration, 1-3µg/ml.

Application

Type

Pep-ELISA, WB-Trf

GALLERY IMAGES





Telephone: (650) 697-3600



