




Telephone: (650) 697-3600

## **GOAT ANTI-DAGLA ANTIBODY**

**SKU:** EB08790

e erest



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 µg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Instructions

Synonym / **Alias** 

KIAA0659|neural stem cell-derived dendrite regulator|diacylglycerol lipase

alpha|NSDDR|DAGLALPHA|DAGL(ALPHA)|C11orf11|diacylglycerol lipase, alpha|DAGLA

**Names Accession** 

NP\_006124.1

**Blocking** 

ID

EBP08790

**Peptide** 

Immunogen

Peptide with sequence C-PAKQDELVISAR, from the C Terminus of the protein sequence according to

NP\_006124.1.

**Peptide** Sequence

C-PAKQDELVISAR

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Shipping

Refrigerated Instructions





Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

**Predicted** 

Human, Mouse, Rat

**Species** Reactive

Human **Species** 

Human

747 **Gene ID** 

Mouse

Gene ID

269060

**Rat Gene ID** 309207 **Product** 

Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_medium.png

IHC Results Paraffin embedded Human Breast, Pancreas and Spleen. Recommended concentration: 5µg/ml.

**ELISA** 

Detection

Antibody detection limit dilution 1:128000.

Limit

Approx 125kDa band observed in Human Liver lysates (calculated MW of 115kDa according to NP\_006124.1). Western

Recommended concentration: 0.3-1µg/ml. Some minor background is detected and is blocked by the Blot

immunizing peptide. Primary incubation was 1 hour.

**Application** 

**Type** 

Pep-ELISA, WB, IHC

## **GALLERY IMAGES**



Telephone: (650) 697-3600



