


GOAT ANTI-DDB1 ANTIBODY

SKU: EB05033



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias Names XPE|damage-specific DNA binding protein 1 (127kD)|DDB p127 subunit|UV-DDB1|XPE-BF|XPCE|XAP1|DDBA|damage-specific DNA binding protein 1, 127kDa|DDB1

Accession ID NP_001914.3

Blocking Peptide EBP05033

Immunogen Peptide with sequence C-DLIKVVEELTRIH, from the C Terminus of the protein sequence according to NP_001914.3.

Peptide Sequence C-DLIKVVEELTRIH

Purification Method Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Shipping Instructions Refrigerated

Predicted Species Human, Mouse, Rat, Dog, Cow

Reactive Species Human, Mouse

| | |
|-------------------------|--|
| Human Gene ID | 1642 |
| Rat Gene ID | 64470 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| IHC Results | Paraffin embedded Human Brain (Cortex). Recommended concentration: 4-6µg/ml. |
| ELISA | |
| Detection Limit | Antibody detection limit dilution 1:32000. |
| Western Blot | Approx 140kDa band observed in lysates of cell line HeLa, HepG2 and Jurkat (calculated MW of 126.9kDa according to Human NP_001914.3). Recommended concentration: 1-2µg/ml. Approx 140kDa band observed in lysates of cell line NIH3T3 and approx 150kDa band observed in lysates of cell line NSO (calculated MW of 126.8kDa according to Mouse NP_056550.1). Recommended concentration: 0.01-0.03µg/ml. Primary incubation was 1 hour. Preliminary testing was unsuccessful on Rat Kidney for this particular batch. |
| Application Type | Pep-ELISA, WB, IHC |

SELECTED REFERENCES

[{"pmid": 19966799, "intro": "**This antibody (previous batch) has been successfully used in WB and IP on Human:**", "title": "A promiscuous alpha-helical motif anchors viral hijackers and substrate receptors to the CUL4-DDB1 ubiquitin ligase machinery.", "author": "Li T, Robert EI, van Breugel PC, Strubin M, Zheng N.", "journal": "Nat Struct Mol Biol. 2010 Jan;17(1):105-11."}]

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

