


Telephone: (650) 697-3600

## **GOAT ANTI-DGAT2 ANTIBODY**



e erest



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

**Alias** 

Names

diacylglycerol O-acyltransferase-like protein 2 isoform 1|diacylglycerol O-acyltransferase homolog 2|GS1999full|HMFN1045|DKFZp686A15125|diacylglycerol O-acyltransferase homolog 2 (mouse)|DGAT2

<strong>Flow Cytometry:</strong> Flow cytometric analysis of HeLa cells. Recommended concentration: Usage

10ug/ml. **Summary** 

**Accession** 

NP\_115953.2; NP\_001240820.1

**Blocking Peptide** 

EBP07661

**Immunogen** 

Peptide with sequence C-KLEHPTQQDIDLYH, from the internal region (near the C Terminus) of the protein sequence according to NP\_115953.2; NP\_001240820.1.









**Product** Comments

This antibody is not expected to cross-react with other DGAT members.

**Peptide** 

C-KLEHPTQQDIDLYH

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method

using the immunizing peptide.

**Shipping** Instructions

Refrigerated

**Predicted** 

**Species** 

Human, Mouse, Rat, Dog

Reactive

**Species** 

Human

Human

84649

Gene ID

Mouse

67800

Gene ID

**Rat Gene ID** 252900

**Product** 

Grade

 $https://prod-vector-labs-pimcore-assets.s 3. us-east-1. a mazon aws.com/assets/products/image/elite\_medium.png$ 

**IHC Results** 

In paraffin embedded Human Liver shows granular cytoplasm staining of hepatocytes. Recommended

concentration: 3µg/ml.

**ELISA** 

**Detection** 

Antibody detection limit dilution 1:128000.

Limit

**Type** 

 $\textbf{Application} \quad \text{Pep-ELISA, IHC, FC}$ 

## **GALLERY IMAGES**

