

## GOAT ANTI-DOCK1 ANTIBODY

**SKU:** EB05183



250kDa  
150kDa  
100kDa  
75kDa  
50kDa  
37kDa  
25kDa

---

## SPECIFICATIONS

<b>Formulation</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
<b>Unit Size</b>	100 µg
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Instructions</b>	
<b>Synonym /</b>	
<b>Alias</b>	ced5 RP11-82L9.1 dedicator of cyto-kinesis 1 DOCK180 DOCK1
<b>Names</b>	
<b>Accession ID</b>	NP_001371.1
<b>Blocking Peptide</b>	EBP05183
<b>Immunogen</b>	Peptide with sequence C-TTRKQTSVDSGIVQ, from the C Terminus of the protein sequence according to NP_001371.1.
<b>Peptide Sequence</b>	C-TTRKQTSVDSGIVQ
<b>Purification Method</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Shipping Instructions</b>	Refrigerated
<b>Predicted Species</b>	Human, Mouse, Rat, Dog, Pig, Cow
<b>Reactive Species</b>	Human
<b>Human Gene ID</b>	1793
<b>Mouse Gene ID</b>	330662
<b>Product Grade</b>	<a href="https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png">https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png</a>
<b>Western Blot</b>	Approx 150-200kDa band seen in A431 cell lysate. Recommended for use at 3-4µg/ml.
<b>Application Type</b>	WB

## DOCUMENTS

- [Data Sheet](#)

## GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa