

Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-EEA1 (AA821-835) ANTIBODY

SKU: EB11689



Telephone: (650) 697-3600



250kDa 150kDa 100kDa 75kDa

50kDa

37kDa

25kDa

20kDa

15kDa



Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Names

Synonym / Alias

zinc finger FYVE domain-containing protein 2|ZFYVE2|MSTP105|MST105|endosome-associated protein p162|EEA1|early endosome-associated protein|early endosome antigen 1, 162kD|early endosome antigen 1

 Immunofluorescence :Strong expression of the protein seen in the vesicles/cytoplasm of Usage

NIH3T3 cells. Recommended concentration: 5µg/ml. Summary

Accession

NP_003557.2

Blocking

ID

EBP11689 **Peptide**

Peptide with sequence C-QETKIQHEELNNRIQ, from the internal region of the protein sequence according to **Immunogen**

NP 003557.2.

Peptide

C-QETKIQHEELNNRIQ Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Human **Species**

Reactive Human, Mouse

Species

Human 8411 Gene ID

Product https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite medium.png

Grade **ELISA**

Detection Antibody detection limit dilution 1:128000.

Limit

Blot

Western

Approx 200kDa band observed in lysates of cell line HEK293 (calculated MW of 162kDa according to

NP_003557.2). This molecular weight is routinely observed by other sources. Recommended concentration:

0.3-1µg/ml. Primary incubatiion 1 hour at room temperature.

Application

Pep-ELISA, WB, IF **Type**

GALLERY IMAGES



Telephone: (650) 697-3600



250kDa 150kDa 100kDa 75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

