
Telephone: (650) 697-3600





## **GOAT ANTI-FBL2 / FBXL2 ANTIBODY**

**SKU:** EB06041



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Storage

Instructions '

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias

 $F-box\ protein\ containing\ leucine-rich\ repeats | DKFZP564P0622 | FBL3 | F-box\ and\ leucine-rich\ repeat\ protein$ 

2|FBL2|FBXL2

Names Usage

<strong>Immunofluorescence:</strong> Strong expression of the protein seen in the cytoplasm of HeLa cells.

Summary

Recommended concentration: 10µg/ml.

Accession ID

NP\_036289.2

Blocking

Peptide EBP06041

Immunogen

Peptide with sequence VFSNNDEGLINKK-C, from the N Terminus of the protein sequence according to

NP\_036289.2.

**Peptide** 

VFSNNDEGLINKK-C

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

**Method** using the immunizing peptide.









**Shipping** Instructions

Refrigerated

**Predicted** 

Human, Rat, Dog, Cow

**Species** Reactive

Human **Species** 

Human

25827 **Gene ID** 

**Product** 

 $https://prod-vector-labs-pimcore-assets.s 3. us-east-1. amazonaws.com/assets/products/image/elite\_medium.png$ Grade

IHC Results Paraffin embedded Human Cortex. Recommended concentration: 8µg/ml.

**ELISA** 

Detection Antibody detection limit dilution 1:128000.

Limit

**Type** 

Western Blot

Preliminary testing showed a band at approx 48kDa in Human Kidney lysate after 0.1µg/ml antibody staining

(calculated MW of 47.1kDa according to NP\_036289.2). Primary incubation 1 hour at room temperature.

Application

Pep-ELISA, IHC, IF

## **GALLERY IMAGES**

