

GOAT ANTI-FBXW2 ANTIBODY

SKU: EB06064

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa

25kDa
20kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	
Alias	F-box and WD repeat domain containing 2 F-box protein Fbw2 Md6 MGC117371 Fwd2 FBW2 F-box and WD-40
Names	domain protein 2 FBXW2
Accession ID	NP_036296.2
Blocking Peptide	EBP06064
Immunogen	Peptide with sequence C-KRGSSFLAGEHPG, from the C Terminus of the protein sequence according to NP_036296.2.
Peptide Sequence	C-KRGSSFLAGEHPG
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Rat, Dog, Pig, Cow
Reactive Species	Human, Mouse, Rat
Human Gene ID	26190
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
IHC Results	In paraffin embedded Human Kidney, shows cytoplasmic staining of epithelial cells in some renal tubules. Recommended concentration, 2-3µg/ml. An anonymous customer found positive results in IHC on Mouse kidney tissue. An anonymous customer found positive results in IHC on Rat kidney tissue.
ELISA Detection Limit	Antibody detection limit dilution 1:128000.
Western Blot	Approx 48kDa band observed in Human Kidney lysates (calculated MW of 48.0kDa according to NP_036296.1). Recommended concentration: 0.3-1.5µg/ml. A stronger band of unknown identity was also consistently observed at appr. 30kDa - both bands were successfully blocked by incubation with the immunising peptide.
Application Type	Pep-ELISA, WB, IHC

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa


150kDa

100kDa

75kDa

50kDa

37kDa



25kDa

20kDa

