

GOAT ANTI-FIBRILLARIN / FBL ANTIBODY

SKU: EB11640

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	rRNA 2'-O-methyltransferase fibrillarin RNU3IP1 RNA, U3 small nucleolar interacting protein
Alias	1 FLRN fibrillarin FIB FBL 34-kD nucleolar scleroderma antigen 34 kDa nucleolar scleroderma antigen
Names	
Accession ID	NP_001427.2
Blocking Peptide	EBP11640
Immunogen	Peptide with sequence C-KRVSISEGDDKIEYR, from the internal region of the protein sequence according to NP_001427.2.
Peptide Sequence	C-KRVSISEGDDKIEYR
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Rat, Dog, Pig, Cow
Reactive Species	Human
Human Gene ID	2091
Mouse Gene ID	14113
Rat Gene ID	292747
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:32000.
Western Blot	Approx 35kDa band observed in lysates of cell line HEK293 (calculated MW of 33.8kDa according to NP_001427.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa