

GOAT ANTI-FOXC1 (INTERNAL) ANTIBODY

SKU: EB07038

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	FOXC1 forkhead box C1 HGNC:3800 ARA FKHL7 FREAC3 IGDA IHG1 IRID1 Forkhead, drosophila, homolog-like
Alias	7 forkhead (Drosophila)-like 7 forkhead-related activator 3 iridogoniodysgenesis type 1 myeloid factor-
Names	delta FREAC-3 RIEG3 forkhead/winged helix-like transcription factor 7
Accession ID	NP_001444.2
Blocking Peptide	EBP07038
Immunogen	Peptide with sequence C-DAVKDKEEKDRLH, from the internal region of the protein sequence according to NP_001444.2.
Peptide Sequence	C-DAVKDKEEKDRLH
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Pig
Reactive Species	Human
Human Gene ID	2296
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA	
Detection Limit	Antibody detection limit dilution 1:32000.
Western Blot	Approx 55kDa band observed in lysates of cell line HEK293 (calculated MW of 56.8kDa according to NP_001444.2). Recommended concentration: 2-4µg/ml.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa