

## GOAT ANTI-HIC2 (AA188-200) ANTIBODY

**SKU:** EB11680

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

---

## SPECIFICATIONS

**Formulation** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 µg

**Storage Instructions** Aliquot and store at -20°C. Minimize freezing and thawing.

**Synonym /**

**Alias** HIC1-related gene on chromosome 22 protein|HIC2|hic-2|hic-3|HRG22|hypermethylated in cancer 2|hypermethylated in cancer 2 protein|ZBTB30|zinc finger and BTB domain-containing protein 30|ZNF907

**Names**

**Accession ID** NP\_055909.2

**Blocking Peptide** EBP11680

**Immunogen** Peptide with sequence C-PQELPQAKGSDE, from the internal region of the protein sequence according to NP\_055909.2.

**Peptide Sequence** C-PQELPQAKGSDE

**Purification Method** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

**Shipping Instructions** Refrigerated

**Predicted Species** Human, Mouse, Rat, Dog, Pig, Cow

**Reactive Species** Human

**Human Gene ID** 23119

**Mouse Gene ID** 58180

**Rat Gene ID** 287940

**Product Grade** [https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\\_medium.png](https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png)

**ELISA**

**Detection Limit** Antibody detection limit dilution 1:128000.

**Western Blot** Approx 60kDa band observed in lysates of cell line HepG2 (calculated MW of 66.2kDa according to NP\_055909.2). Recommended concentration: 0.1-0.3µg/ml.

**Application Type** Pep-ELISA, WB

## DOCUMENTS

- [Data Sheet](#)

## GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

