

# GOAT ANTI-HISTAMINE RECEPTOR H1(INTERNAL) ANTIBODY

**SKU:** EB07699

250kDa  
150kDa  
100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

## SPECIFICATIONS

<b>Formulation</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
<b>Unit Size</b>	100 µg
<b>Storage Instructions</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Synonym / Alias Names</b>	histamine receptor H1 H1-R HRH1 hisH1 histamine receptor, subclass H1
<b>Accession ID</b>	NP_000852.1
<b>Blocking Peptide</b>	EBP07699
<b>Immunogen</b>	Peptide with sequence QQTSVRREDKCETD, from the internal region of the protein sequence according to NP_000852.1.
<b>Peptide Sequence</b>	QQTSVRREDKCETD
<b>Purification Method</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Shipping Instructions</b>	Refrigerated
<b>Predicted Species</b>	Human
<b>Reactive Species</b>	Human
<b>Human Gene ID</b>	3269
<b>Product Grade</b>	<a href="https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png">https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png</a>
<b>IHC Results</b>	In paraffin embedded Human Lung shows membranous staining of epithelial cells in tertiary bronchus and in alveoli. Recommended concentration, 2-4µg/ml.
<b>ELISA Detection Limit</b>	Antibody detection limit dilution 1:16000.
<b>Western Blot</b>	Approx 55kDa band observed in Human Heart lysates (calculated MW of 55.8kDa according to NP_000852.1). Recommended concentration: 0.5-1.5µg/ml.
<b>Application Type</b>	Pep-ELISA, WB, IHC

## DOCUMENTS

- [Data Sheet](#)

## GALLERY IMAGES

