


Telephone: (650) 697-3600

## **GOAT ANTI-HOXD13 ANTIBODY**

**SKU:** EB09834



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

HOXD13|SPD|HOX4I|homeobox D13|homeo box D13|homeo box 4I|BDSD|BDE Alias

Names

Usage <strong>Immunofluorescence:</strong> Strong expression of the protein seen in the nucleus of HEK293 and U2OS

cells. Recommended concentration: 10µg/ml. Summary

Accession NP 000514.2

**Blocking** 

EBP09834 Peptide

Peptide with sequence C-HGVGLQQNALKSSPH, from the internal region of the protein sequence according to **Immunogen** 

NP 000514.2.

**Peptide** C-HGVGLQQNALKSSPH Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

**Shipping** 

Refrigerated Instructions









**Predicted** 

Human, Mouse, Rat, Cow, Pig

**Species** Reactive

Human, Rat **Species** 

Human

3239

**Gene ID** 

Mouse

15433 Gene ID

**Rat Gene ID** 288154

**Product** 

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_plus\_medium.png Grade

IHC Results Paraffin embedded Human Prostate. Recommended concentration: 8µg/ml.

**ELISA** 

Detection Antibody detection limit dilution 1:128000.

Limit

Blot

Western Approx 37kDa band observed in Rat Brain lysates (calculated MW of 34.9kDa according to NP\_001099356.1).

Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

Type

**Application** Pep-ELISA, WB, IF, IHC

## **GALLERY IMAGES**

