



Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-LAT2 ANTIBODY

SKU: EB12057



Email: customerservice@vectorlabs.com

Telephone: <u>(650)</u> 697-3600

250kDa 150kDa 100kDa

75kDa


50kDa

37kDa

25kDa

20kDa

15kDa





Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias

Names

non-T-cell activation linker|membrane-associated adapter molecule|linker for activation of T-cells family member 2|linker for activation of T cells, transmembrane adaptor 2|linker for activation of B-cells|Williams-Beuren syndrome

chromosomal region 5 protein|Williams-Beuren syndrome chromosomal region 15

protein|WSCR5|WBSCR5|WBSCR15|NTAL|LAB|linker for activation of T cells family, member 2|LAT2

Immunofluorescence: Strong expression of the protein seen in U937 cells. Recommended Usage concentration: 10μg/ml. Flow Cytometry: Flow cytometric analysis of U937 cells. Summary

Recommended concentration: 10ug/ml.

Accession

ID

NP 054865.2

Blocking Peptide

EBP12057

Immunogen

Peptide with sequence C-RSEKIYQQRSLRED, from the internal region of the protein sequence according to

NP 054865.2.

Product Comments

Reported variants represent identical protein: NP_115853.2, NP_115852.1, NP_054865.2

Peptide

C-RSEKIYOORSLRED Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Species

Human

Reactive

Species

Human

Human **Gene ID**

7462

Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_plus_medium.png

Grade **ELISA**

Detection

Antibody detection limit dilution 1:128000.

Limit

Western

Approx. 26kDa band observed in Human Peripheral Blood Monocytes lysates and Human Spleen, and approx. 30kDa in lysates of cell lines Daudi and U937 (calculated MW of 26.6kDa according to NP_054865.2). Recommended

concentration: 0.1-2µg/ml. Primary incubation 1 hour at room temperature.

Application

Type

Blot

Pep-ELISA, WB, IF, FC





Telephone: (650) 697-3600

















25kDa

20kDa

15kDa





Telephone: (650) 697-3600



