
Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

## **GOAT ANTI-MYD88 ANTIBODY**





## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias

myeloid differentiation primary response gene 88|MYD88D|myeloid differentiation primary response gene

(88)|MYD88

**Names** Usage

<strong>Immunofluorescence:</strong> Strong expression of the protein seen in U2OS and Jurkat cells.

Summary

Recommended concentration: 10µg/ml. <strong>Flow Cytometry:</strong> Flow cytometric analysis of Jurkat

cells. Recommended concentration: 10ug/ml.

Accession ID

NP 001166038.2; NP 002459.3; NP 001166039..2; NP 001361717.1

**Blocking** Peptide

EBP06667


**Immunogen** 

Peptide with sequence C-IKYKAMKKEFP, from the internal region of the protein sequence according to  $NP\_001166038.2; \ NP\_002459.3; \ NP\_001166039..2; \ NP\_001361717.1.$ 











This antibody is expected to recognize reported isoforms 1, 2, 3 and 9.

**Comments Peptide** 

e erest

C-IKYKAMKKEFP

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

**Shipping** 

Refrigerated Instructions

**Predicted Species** 

Human, Mouse, Rat, Dog

Reactive

Human **Species** 

Human

4615

**Gene ID** 

Mouse 17874

Gene ID

**Rat Gene ID** 301059

**Product** 

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_plus\_medium.png Grade

IHC Results Paraffin embedded Human Tonsil. Recommended concentration: 4-6µg/ml.

**ELISA** 

**Detection** Antibody detection limit dilution 1:128000.

Limit

Blot

Western Approx 28kDa band observed in Human Spleen lysates (calculated MW of 28.3kDa according to NP\_001166039.2).


Recommended concentration: 0.3-0.5µg/ml. Primary incubation1 hour at room temperature.


Application


Type

Pep-ELISA, WB, IHC, IF, FC

## **GALLERY IMAGES**











Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

