

GOAT ANTI-NOXA ANTIBODY

SKU: EB09119

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	adult T cell leukemia-derived PMA-responsive NOXA APR phorbol-12-myristate-13-acetate-induced protein
Alias	1 PMAIP1
Names	
Usage	Flow Cytometry: Flow cytometric analysis of LNCaP cells. Recommended concentration:
Summary	10ug/ml.
Accession ID	NP_066950.1; NP_001369544.1
Blocking Peptide	EBP09119
Immunogen	Peptide with sequence C-TQLRRFGDKLNFRQK, from the internal region of the protein sequence according to NP_066950.1; NP_001369544.1.
Peptide Sequence	C-TQLRRFGDKLNFRQK
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human
Reactive Species	Human
Human Gene ID	5366
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:16000.
Western Blot	Approx 16kDa band observed in Human Bone Marrow lysates and in lysates of cell line LNCaP and in preliminary testing of Human Cerebellum lysate and Peripheral blood monocyte cell lysate (calculated MW of 6.03kDa according to NP_066950.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.3-1ug/ml. Primary incubation 1 hour at room temperature.
Application Type	Pep-ELISA, WB, FC

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa

