


Telephone: (650) 697-3600

## **GOAT ANTI-NUDC ANTIBODY**

**SKU:** EB09516



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Storage Instructions

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias

 $nuclear\ distribution\ gene\ C\ homolog|OTTHUMP00000004405|NPD011|MNUDC|HNUDC|nuclear\ distribution\ gene$ 

C homolog (A. nidulans)|NUDC

Usage Summary

**Names** 

<strong>Immunofluorescence:</strong> Strong expression of the protein seen in the cytoplasm of A431 cells.

Recommended concentration: 10µg/ml.

Accession

NP\_006591.1

Blocking

**Peptide** 

EBP09516

**Immunogen** 

Peptide with sequence C-PNLGNGADLPNYR, from the internal region of the protein sequence according to

NP\_006591.1.









Peptide C-PN

C-PNLGNGADLPNYR

Sequence

**Purification** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

**Method** using the immunizing peptide.

Shipping

Instructions Re

Refrigerated

Predicted Species

Human, Mouse, Rat, Dog, Cow

Reactive

Species Human, Rat

Human

Gene ID

Mouse

Gene ID

Rat Gene ID 29648

**Product** 

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_medium.png

Grade ELISA

**Detection** 

Antibody detection limit dilution 1:32000.

Limit

Western Blot Approx 40kDa band observed in lysates of cell lines HeLa. HepG2 and K562, and approx. 38kDa in lysates of cell line KNRK (calculated MW of 38.2kDa according to Human NP\_006591.1) and 38.4kDa according to Rat NP\_058967.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

Application

Type

Pep-ELISA, WB, IF

## **GALLERY IMAGES**

