

GOAT ANTI-OXER1 (AA389-400) ANTIBODY

SKU: EB10894

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

| | |
|------------------------------|---|
| Formulation | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. |
| Unit Size | 100 µg |
| Storage | Aliquot and store at -20°C. Minimize freezing and thawing. |
| Instructions | |
| Synonym / | oxoeicosano G-protein coupled receptor R527 G-protein coupled receptor 170 5-oxo-ETE acid G-protein- |
| Alias | coupled receptor 1 5-oxo-ETE G-protein coupled receptor TG1019 GPR170 GPCR oxoeicosanoid (OXE) |
| Names | receptor 1 OXER1 |
| Accession ID | NP_683765.1 |
| Blocking Peptide | EBP10894 |
| Immunogen | Peptide with sequence C-SRQWRYREASRK, from the C Terminus of the protein sequence according to NP_683765.1. |
| Peptide Sequence | C-SRQWRYREASRK |
| Purification Method | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Shipping Instructions | Refrigerated |
| Predicted Species | Human |
| Reactive Species | Human |
| Human Gene ID | 165140 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| ELISA | |
| Detection Limit | Antibody detection limit dilution 1:16000. |
| Western Blot | Approx 38kDa band observed in Human Lung lysates (calculated MW of 45.8kDa according to NP_683765.1). This molecular weight is routinely observed by other sources. Recommended concentration: 1-3µg/ml. |
| Application Type | Pep-ELISA, WB |

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa