

GOAT ANTI-PCQAP / MED15 ANTIBODY

SKU: EB09173

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias Names MED15|mediator complex subunit
15|ARC105|CAG7A|CTG7A|DKFZp686A2214|DKFZp762B1216|FLJ42282|FLJ42935|PCQAP|TIG-1|TIG1|TNRC7|CTG repeat protein 7a|PC2 (positive cofactor 2, multiprotein complex) glutamine/Q-rich-associated protein|PC2-glutamine-rich-associated protein|TPA inducible gene-1|TPA inducible protein|activator-recruited cofactor, 105-kD|positive cofactor 2, glutamine/Q-rich-associated protein|trinucleotide repeat containing 7

Accession ID NP_001003891.1; NP_056973.2

Blocking Peptide EBP09173

Immunogen Peptide with sequence C-PPQSQTQPLVSQAQ, from the internal region of the protein sequence according to NP_001003891.1; NP_056973.2.

Product Comments This antibody is expected to recognize both reported isoforms (NP_001003891.1; NP_056973.2).

Peptide Sequence C-PPQSQTQPLVSQAQ

Purification Method Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Shipping Instructions Refrigerated

Predicted Species Human, Mouse

Reactive Species Human

Human Gene ID 51586

Mouse Gene ID 94112

Product Grade https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png

ELISA Detection Limit Antibody detection limit dilution 1:8000.

Western Blot Approx. 90kDa band observed in Human Lung lysates (calculated MW of 86.8kDa according to NP_001003891.1). Recommended concentration: 1-3µg/ml.

Application Type Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

