
Telephone: (650) 697-3600



Email: customerservice@vectorlabs.com

GOAT ANTI-PHLDA2 / IPL ANTIBODY

SKU: EB06184







Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size

Aliquot and store at -20°C. Minimize freezing and thawing.

tumor-supressing STF cDNA 3|tumor suppressing subtransferable candidate 3|tumor suppressing subchromosomal

transferable fragment cDNA 3|pleckstrin homology-like domain family A member 2|p17-Beckwith-Wiedemann Synonym / region 1C|imprinted in placenta and liver|tumor-supressing STF cDNA 3|tumor suppressing subtransferable Alias candidate 3|PHLDA2|HLDA2|BWR1C|BRW1C|IPL|TSSC3|pleckstrin homology-like domain, family A, member Names

2|PHLDA2

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of HeLa cells. Usage

Recommended concentration: 10µg/ml. Flow Cytometry: Flow cytometric analysis of MCF7 **Summary**

cells. Recommended concentration: 10ug/ml.

Accession

NP_003302.1

Blocking

ID

EBP06184 **Peptide**

Immunogen | NP_003302.1. Peptide with sequence C-EPSRPSPQPKPRTP, from the C Terminus of the protein sequence according to

Peptide

C-EPSRPSPQPKPRTP Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Human **Species**

Reactive Human **Species**

Human

7262 **Gene ID**

Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_plus_medium.png Grade

IHC Results Paraffin embedded Human Placenta and Prostate. Recommended concentration: 3.75µg/ml.

ELISA

Detection Antibody detection limit dilution 1:32000.

Limit

Blot

Type

Western

Approx. 22kDa band observed in lysates of cell line MDA-MB-231 and in preliminary testing of MCF7 cell lysate (calculated MW of 17.1kDa according to NP_003302.1). This molecular weight has been observed by other sources.

Recommended concentration 2-3 μ g/ml Primary incubation 1 hour at room temperature.


Application


Pep-ELISA, WB, IF, IHC, FC


GALLERY IMAGES











50kDa - 37kDa •