



Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-PICK1 ANTIBODY

SKU: EB07467



Telephone: (650) 697-3600



250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

25kDa 20kDa

15kDa

10kDa





Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias Names protein kinase C, alpha binding protein|protein interacting with C kinase 1|PRKCA-binding protein|OTTHUMP00000028509|PRKCABP|MGC15204|protein interacting with PRKCA 1|PICK1

Accession

NP_036539.1

Blocking

Peptide

EBP07467

Peptide with sequence C-HYNKLQADPKQ, from the internal region of the protein sequence according to **Immunogen**

NP 036539.1.

Product Comments

Reported variants represent identical protein (NP 036539.1; NP 001034672.1; NP 001034673.1).

Peptide

Sequence

C-HYNKLQADPKQ

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Human, Mouse

9463

Shipping

Refrigerated Instructions

Predicted

Species

Human, Mouse, Rat, Dog, Pig, Cow

Reactive **Species**

Human

Gene ID

Mouse

18693 Gene ID

Rat Gene ID 84591

Product Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png

IHC Results Paraffin embedded Human Colon and Prostate. Recommended concentration: 5µg/ml.

ELISA

Detection Antibody detection limit dilution 1:16000.

Limit

Western

Approx 50kDa band observed in Mouse Brain lysates (calculated MW of 46.6kDa according to NP 036539.1, NP_001034672.1and NP_001034673.1). Recommended concentration: 1-3μg/ml. Primary incubation was 1

Blot

Application

Pep-ELISA, WB, IHC

Type



Telephone: (650) 697-3600

GALLERY IMAGES

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa



