

GOAT ANTI-PLA2G4C ANTIBODY

SKU: EB10114

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	PLA2G4C phospholipase A2, group IVC (cytosolic, calcium-independent) phospholipase A2, group IVC
Alias	IVC phospholipase A2, cytosolic, calcium-independent, gamma FLJ44164 FLJ42247 DKFZp586C0423 cytosolic
Names	phospholipase A2 gamma CPLA2-gamma
Accession ID	NP_003697.2; NP_001152794.1; NP_001152795.1
Blocking Peptide	EBP10114
Immunogen	Peptide with sequence C-NTEVIREYIFDQLR, from the internal region of the protein sequence according to NP_003697.2; NP_001152794.1; NP_001152795.1.
Product Comments	This antibody is expected to recognize all reported isoforms (NP_003697.2; NP_001152794.1; NP_001152795.1).
Peptide Sequence	C-NTEVIREYIFDQLR
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human
Reactive Species	Human
Human Gene ID	8605
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:128000.
Western Blot	Approx 65kDa band observed in Human Brain (Hippocampus), Liver, Pancreas, and Spleen lysates (calculated MW of 62.5kDa according to NP_001152794.1). Recommended concentration: 0.03-0.1µg/ml.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

Goat Anti-PLA2G4C Antibody

<https://everestbiotech.com/products/goat-anti-pla2g4c-antibody/>

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa