


GOAT ANTI-POFUT1 ANTIBODY

SKU: EB11528



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

| | |
|------------------------------|--|
| Unit Size | 100 µg |
| Storage Instructions | Aliquot and store at -20°C. Minimize freezing and thawing. |
| Synonym / Alias | protein O-fucosyltransferase 1 POFUT1 peptide-O-fucosyltransferase 1 OTTHUMP00000030584 OTTHUMP00000030583 O-FUT O-FucT-1 O-Fuc-T o-fucosyltransferase |
| Names | protein MGC2482 KIAA0180 GDP-fucose protein O-fucosyltransferase 1 FUT12 |
| Usage Summary | Purchase the +ve control! The transfected lysate used in QC is available for sale. Please email sales@everestbiotech.com to request datasheet, pricing and delivery information. |
| Accession ID | NP_056167.1; NP_758436.1 |
| Blocking Peptide | EBP11528 |
| Immunogen | Peptide with sequence C-TNLHVSQYQYFKLEP, from the internal region of the protein sequence according to NP_056167.1; NP_758436.1. |
| Product Comments | This antibody is expected to recognize both reported isoforms (NP_056167.1; NP_758436.1). |
| Peptide Sequence | C-TNLHVSQYQYFKLEP |
| Purification Method | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Shipping Instructions | Refrigerated |
| Predicted Species | Human, Mouse, Rat, Dog, Pig, Cow |
| Reactive Species | Human |
| Human Gene ID | 23509 |
| Mouse Gene ID | 140484 |
| Rat Gene ID | 311551 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| IHC Results | In paraffin embedded Human Cerebral Cortex shows staining of the cytoplasm in neuronal bodies. Recommended concentration, 2-4µg/ml. |
| ELISA Detection Limit | Antibody detection limit dilution 1:16000. |
| Western Blot | In transfected HEK293 transiently expressing Human POFUT1 (myc and DYKDDDDK tagged), a band of approx. 48kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc antibody. Recommended concentration, 0.3-1µg/ml. |
| Application Type | Pep-ELISA, WB-Trf, IHC |

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

