


Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-POLO-LIKE KINASE 3 (AA320-333) ANTIBODY

SKU: EB12012



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias Names serine/threonine-protein kinase PLK3|proliferation-related kinase|cytokine-inducible serine/threonine-protein kinase|cytokine-inducible kinase|PLK-3|FGF-inducible kinase|PRK|FNK|CNK|polo-like kinase 3|PLK3

Accession

NP 004064.2

Blocking

Peptide

EBP12012

Immunogen

Peptide with sequence PDRLPISSCVTVPD, from the internal region of the protein sequence according to NP 004064.2.

Peptide Sequence

PDRLPISSCVTVPD

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method

using the immunizing peptide.





Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

Shipping Instructions

Refrigerated

Predicted

Human

Species Reactive

Human

Species

Human **Gene ID**

1263

Product

 $https://prod-vector-labs-pimcore-assets.s 3. us-east-1. a mazon aws.com/assets/products/image/elite_medium.png$ Grade

IHC Results

In paraffin embedded Human Kidney shows vesicular staining in the cystoplasm of collective tubules.

Recommended concentration, 5-10µg/ml.

ELISA

Detection

Antibody detection limit dilution 1:128000.

Limit

Western

Approx 70kDa band observed in lysates of cell line HeLa (calculated MW of 71.6kDa according to NP_004064.2). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at rroom temperature. **Blot**

Application

Type


Pep-ELISA, WB, IHC

GALLERY IMAGES



Telephone: (650) 697-3600





250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa