

GOAT ANTI-PRDM10 ANTIBODY

SKU: EB05717



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias Names PRDM zinc finger transcription factor|PR-domain family member 7|PFM7|MGC131802|tristanin|PR domain containing 10|KIAA1231|PRDM10

Accession ID

NP_064613.2; NP_955469.1; NP_955470.1; NP_955471.1; NP_001354824.1; NP_001354826.1

Blocking Peptide

EBP05717

Immunogen Peptide with sequence C-NGNGSSEVHITKP, from the C Terminus of the protein sequence according to NP_064613.2; NP_955469.1; NP_955470.1; NP_955471.1; NP_001354824.1; NP_001354826.1.

Peptide Sequence C-NGNGSSEVHITKP

Purification Method Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Shipping Instructions Refrigerated

Predicted Species Human, Cow

Human Gene ID 56980

Product Grade https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/aspiring_medium.png

ELISA

Detection Limit Antibody detection limit dilution 1:32000.

Western Blot No signal obtained yet but low background observed in Human Brain and Human Kidney lysates at up to 1µg/ml.
We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates?

Application Type Pep-ELISA

DOCUMENTS

- [Data Sheet](#)