




GOAT ANTI-PRKAB2 ANTIBODY

SKU: EB08470



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias **Names**

AMPK beta-2 chain|AMPK beta 2|AMP-activated protein kinase beta 2 non-catalytic subunit|5\1-AMP-activated protein kinase, beta-2 subunit|MGC61468|protein kinase, AMP-activated, beta 2 non-catalytic subunit|PRKAB2

Usage **Summary**

Immunofluorescence: Strong expression of the protein seen in the nuclei of MCF7 and A431 cells. Recommended concentration: 10µg/ml. Flow Cytometry: Flow cytometric analysis of

A431 cells. Recommended concentration: 10ug/ml.

Accession

NP_005390.1

Blocking Peptide

EBP08470

Immunogen NP_005390.1. Peptide with sequence C-PYGQEMYAFRSE, from the internal region of the protein sequence according to









Peptide C-PYGQEMYAFRSE

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Human, Mouse, Rat, Cow

Species Reactive

Human Species

Human

5565 **Gene ID**

Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_plus_medium.png

Grade

IHC Results Paraffin embedded Human Liver and Small Intestine. Recommended concentration: 3.75µg/ml.

ELISA

Detection Antibody detection limit dilution 1:8000.

Limit

Application **Type**

Pep-ELISA, IF, IHC, FC

GALLERY IMAGES

