

# GOAT ANTI-PU.1 ANTIBODY

**SKU:** EB08429



250kDa  
150kDa  
100kDa  
75kDa  
50kDa  
37kDa  
25kDa  
20kDa  
15kDa

## SPECIFICATIONS

**Formulation** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 µg

**Storage Instructions** Aliquot and store at -20°C. Minimize freezing and thawing.

**Synonym / Alias Names** hematopoietic transcription factor PU.1|SPI-1 proto-oncogene|31 kDa transforming protein|SPI-A|SPI-1|SFPI1|hCG\_25181|spleen focus forming virus (SFFV) proviral integration oncogene spi1|SPI1|PU.1

**Accession ID** NP\_001074016.1; NP\_003111.2

**Blocking Peptide** EBP08429

**Immunogen** Peptide with sequence C-DLYQRQTHEYY, from the internal region of the protein sequence according to NP\_001074016.1; NP\_003111.2.

**Product Comments** This antibody is expected to recognise both reported isoforms (NP\_001074016.1; NP\_003111.2).

**Peptide Sequence** C-DLYQRQTHEYY

**Purification Method** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

**Shipping Instructions** Refrigerated

**Predicted Species** Human, Rat

**Reactive Species** Human

**Human Gene ID** 6688

**Rat Gene ID** 366126

**Product Grade** [https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\\_medium.png](https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png)

**IHC Results** Paraffin embedded Human Spleen and Colon (lymphoid aggregate). Recommended concentration: 3.75µg/ml.

**ELISA Detection Limit** Antibody detection limit dilution 1:32000.

**Western Blot** Approx 40kDa band observed in lysates of DAUDI and K562 (calculated MW of 31.2kDa according to NP\_001074016.1 and of 31.1kDa according to NP\_003111.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Harendza S et al, J Biol Chem. 2000 Jun 30;275(26):19552-9.; PMID: 10867017). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour.

**Application Type** Pep-ELISA, WB, IHC

## DOCUMENTS

- [Data Sheet](#)

## GALLERY IMAGES

