
Telephone: (650) 697-3600





GOAT ANTI-SGLT1 ANTIBODY

SKU: EB09310



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μα

Storage

Aliquot and store at -20°C. Minimize freezing and thawing. Instructions

Synonym / **Alias**

solute carrier family 5 (sodium/glucose transporter), member 1|sodium/glucose cotransporter 1|Human Na+/glucose cotransporter 1 mRNA, complete cds|SGLT1|NAGT|D22S675|solute carrier family 5

Names (sodium/glucose cotransporter), member 1|SLC5A1

Accession

NP_000334.1; NP_001243243.1 ID

Blocking Peptide

EBP09310

Immunogen

Peptide with sequence C-KETIEIETQVPEKKK, from the internal region of the protein sequence according to

NP_000334.1; NP_001243243.1.

Product Comments

This antibody is expected to recognize isoforms 1 ((NP_000334.1) and isoform 2 (NP_001243243.1).

Peptide Sequence

C-KETIEIETQVPEKKK









Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Shipping Instructions

Refrigerated

Predicted

Human **Species**

Reactive

Human Species

Human

6523 Gene ID

Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png Grade

IHC Results Paraffin embedded Human Small Intestine. Recommended concentration: 5µg/ml.

ELISA

Detection Antibody detection limit dilution 1:32000.

Limit

Preliminary testing showed a band at approx. 60kDa in Human Gall Bladder lysate and in Jurkat cell lysate Western after 1-3µg/ml antibody staining (calculated MW of 60.1kDa according to NP 001243243.1). Primary incubation Blot

1 hour at room temperature.

Application

Pep-ELISA, IHC **Type**

SELECTED REFERENCES

[{"pmid": 34198013, "intro": "This antibody (previous batch) has been successfully used in Western blot and IHC on Mouse:", "title": "Aberrant Epithelial Differentiation Contributes to Pathogenesis in a Murine Model of Congenital Tufting Enteropathy.", "author": "Barun Das, Kevin Okamoto, John Rabalais, Jocelyn A. Young, Kim E. Barrett, and Mamata Sivagnanam", "journal": "Cell Mol Gastroenterol Hepatol. 2021; 12(4): 1353-1371."}, {"pmid": 28174043, "intro": "This antibody (previous batch) has been successfully used in Western blot on Mouse:", "title": "Sodium influx through cerebral sodiumglucose transporter type 1 exacerbates the development of cerebral ischemic neuronal damage. ", "author": "Yamazaki Y, Harada S, Wada T, Hagiwara T, Yoshida S, Tokuyama S", "journal": "Eur J Pharmacol. 2017 Mar 15;799:103-110."}, {"pmid": 30503674, "intro": "This antibody (previous batch) has been successfully used in Western blot on Mouse:", "title":

"Activation of c-Jun N-terminal kinase and p38 after cerebral ischemia upregulates cerebral sodium-glucose transporter type 1", "author": "Yui Yamazaki, Kyoko Arita, Shinichi Harada, Shogo Tokuyama", "journal": "Journal of Pharmacological Sciences (November 2018)"}]

GALLERY IMAGES









