

Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

## GOAT ANTI-SH2D1A/SLAM ASSOCIATED PROTEIN ANTIBODY

**SKU:** EB07013



Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

250kDa 150kDa 100kDa

75kDa

50kDa


37kDa

25kDa

20kDa

15kDa

10kDa





Telephone: (650) 697-3600

## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

T cell signal transduction molecule SAP; signaling lymphocyte activation molecule-associated protein SLAM-

associated protein|Duncan's disease|Duncan's disease (lymphoproliferative

Synonym /

syndrome)|FLJ92177|FLJ18687|RP5-1052M9.3|T cell signal transduction molecule SAP|SLAM (signaling

**Alias** lymphocyte activation molecule) associated protein|SH2 domain protein **Names** 

1A|XLPD|XLP|SAP|MTCP1|LYP|IMD5|EBVS|DSHP|HGNC:10820|SH2 domain protein 1A, Duncan\1s disease

(lymphoproliferative syndrome)|SH2D1A

**Usage** Summary <strong>Immunoprecipitation:</strong>This antibody was deemed fit for IP under native conditions

(observations from anonymous customer).

Accession

ID

NP\_002342.1; NP\_001108409.1

**Blocking** 

**Peptide** 

EBP07013

Peptide with sequence C-QYPVEKKSSARSTQ, from the internal region of the protein sequence according to **Immunogen** 

NP 002342.1; NP 001108409.1.

**Peptide** 

Sequence

C-QYPVEKKSSARSTQ

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

using the immunizing peptide. Method

Shipping

Refrigerated Instructions

**Predicted** Species

Human, Pig

Reactive

**Species** 

Human

Human

4068 Gene ID

**Product** 

**Grade** 

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_medium.png

**ELISA** 

**Detection** 

Antibody detection limit dilution 1:32000.

Limit

Western

Approx 16kDa band observed in Human Lymph Node lysates (calculated MW of 14.2kDa according to


Blot NP\_002342.1).. Recommended concentration: 0.2-0.6μg/ml.

**Application** 

Pep-ELISA, WB, IP

**Type** 







## **GALLERY IMAGES**

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa