

GOAT ANTI-SH2D1A/SLAM ASSOCIATED PROTEIN ANTIBODY

SKU: EB07013



250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa
10kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym / Alias Names	T cell signal transduction molecule SAP; signaling lymphocyte activation molecule-associated protein SLAM-associated protein Duncan's disease Duncan's disease (lymphoproliferative syndrome) FLJ92177 FLJ18687 RP5-1052M9.3 T cell signal transduction molecule SAP SLAM (signaling lymphocyte activation molecule) associated protein SH2 domain protein 1A XLPD XLP SAP MTCP1 LYP IMD5 EBVS DSHP HGNC:10820 SH2 domain protein 1A, Duncan\1s disease (lymphoproliferative syndrome) SH2D1A
Usage Summary	Immunoprecipitation: This antibody was deemed fit for IP under native conditions (observations from anonymous customer).
Accession ID	NP_002342.1; NP_001108409.1
Blocking Peptide	EBP07013
Immunogen	Peptide with sequence C-QYPVEKKSSARSTQ, from the internal region of the protein sequence according to NP_002342.1; NP_001108409.1.
Peptide Sequence	C-QYPVEKKSSARSTQ
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Pig
Reactive Species	Human
Human Gene ID	4068
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:32000.
Western Blot	Approx 16kDa band observed in Human Lymph Node lysates (calculated MW of 14.2kDa according to NP_002342.1).. Recommended concentration: 0.2-0.6µg/ml.
Application Type	Pep-ELISA, WB, IP

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

