



Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-SLY ANTIBODY

SKU: EB05364

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

Alias likely ortholog of mouse SH3 gene SLY|753P9|SLY|chromosome X open reading frame 9|CXorf9

Names

Please note that the band size we observe is higher than the predicted 42kDa. We do not have explanation for this at present. However, CXorf9, has a consensus glycosylation site. Glycosylation can affect migration of proteins on

Usage **Summary** SDS-PAGE [Hames, B.D and Rickwood, D., eds. Gel electrophoresis of proteins: a practical approach. Oxford

University Press].

Accession

NM 018990 ID

Immunogen Peptide with sequence C-EQLQGLSLAGAP, from the C Terminus of the protein sequence according to NM_018990.

Peptide

C-EQLQGLSLAGAP Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Human **Species**

Reactive

Human **Species**

Human

54440 Gene ID

Product

Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/aspiring_medium.png

Western

Western blot: Approx 55-60kDa band seen in Daudi lysate. Recommended for use at 1-3µg/ml. Blot