

GOAT ANTI-SMAD9 ANTIBODY

SKU: EB07200

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym / Alias	Mothers against decapentaplegic, drosophila, homolog of, 9 MAD, mothers against decapentaplegic homolog 9 (Drosophila) MAD, mothers against decapentaplegic homolog 9 MAD (mothers against decapentaplegic, Drosophila) homolog 9 SMAD8B SMAD8A MADH9 MADH6 HGNC:6774 SMAD, mothers against DPP homolog 9 (Drosophila)] SMAD9
Accession ID	NP_005896.1
Blocking Peptide	EBP07200
Immunogen	Peptide with sequence C-HSEPLMPHNATYPD, from the internal region of the protein sequence according to NP_005896.1.
Product Comments	This antibody is not expected to cross-react with SMAD1 and SMAD5.
Peptide Sequence	C-HSEPLMPHNATYPD
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Rat, Dog
Reactive Species	Human
Human Gene ID	4093
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
IHC Results	Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.
ELISA Detection Limit	Antibody detection limit dilution 1:64000.
Western Blot	Approx 50kDa band observed in Human Skin lysates (calculated MW of 48.6kDa according to NP_005896.1). Recommended concentration: 0.3-1.0 µg/ml. Primary incubation was 1 hour.
Application Type	Pep-ELISA, WB, IHC

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa

