

GOAT ANTI-SOCS7 ANTIBODY

SKU: EB08929

250kDa
150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

| | |
|------------------------------|---|
| Formulation | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. |
| Unit Size | 100 µg |
| Storage | Aliquot and store at -20°C. Minimize freezing and thawing. |
| Instructions | |
| Synonym / | |
| Alias | SOCS7 suppressor of cytokine signaling 7 NAP4 Nck, Ash and phospholipase C binding protein |
| Names | |
| Accession ID | NP_055413.1 |
| Blocking Peptide | EBP08929 |
| Immunogen | Peptide with sequence C-DAGRKPKLTRTQS, from the internal region of the protein sequence according to NP_055413.1. |
| Peptide Sequence | C-DAGRKPKLTRTQS |
| Purification Method | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Shipping Instructions | Refrigerated |
| Predicted Species | Human |
| Reactive Species | Human |
| Human Gene ID | 30837 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| ELISA Detection Limit | Antibody detection limit dilution 1:32000. |
| Western Blot | Approx. 60kDa band observed in lysates of Daudi (calculated MW of 63.0kDa according to NP_055413.1). Recommended concentration: 1-3µg/ml. An additional band of unknown identity was also consistently observed at 28kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? |
| Application Type | Pep-ELISA, WB |

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa