

Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

## **GOAT ANTI-TCF3 / ITF1, BIOTINYLATED ANTIBODY**

**SKU:** EB07587-B





Telephone: (650) 697-3600

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa



Telephone: (650) 697-3600

## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 100 μg

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

transcription factor E2-alpha|MGC129648|MGC129647|E2A immunoglobulin enhancer-binding factor

Synonym /

E12/E47|vitamin D receptor-interacting repressor|transcription factor ITF-1|transcription factor 3 variant 3|transcription factor 3 (E2A immunoglobulin enhancer binding factors E12/E47)|negative vitamin D response element-binding protein|kappa-E2-binding factor|immunoglobulin transcription factor 1|helix-loop-helix protein

HE47|class B basic helix-loop-helix protein 21|VDR interacting repressor|NOL1-TCF3

fusion|bHLHb21|VDIR|TCF-3|ITF1|E47|E2A|transcription factor 3|TCF3

Accession

**Alias** 

**Names** 

NP 003191.1

**Blocking** 

**Peptide** 

EBP07587-B

**Immunogen** 

Peptide with sequence C-KAPRARTSPDEDED., from the internal region of the protein sequence according to

NP 003191.1.

**Product** 

Comments

This antibody is expected to recognize reported isoform E12 (NP\_003191.1) only.

**Peptide** 

Sequence

C-KAPRARTSPDEDED.

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method

using the immunizing peptide.

Shipping

Instructions

Refrigerated

**Predicted** 

Human Species

Reactive **Species** 

Human

Human

6929

Gene ID

**Product Grade** 

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_medium.png

**ELISA** 

**Detection** 

Western

Antibody detection limit dilution 1:128000.

Limit

**Blot** 

Approx 75kDa band observed in lysates of cell line Daudi (calculated MW of 67.6kDa according to NP\_003191.1). See non-biotinylated parental product's datasheet for further QC data. Recommended

concentration: 0.3-1µg/ml.

**Application** 

**Type** 

Pep-ELISA, WB







## **GALLERY IMAGES**

250kDa 150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa