






GOAT ANTI-TIM3 / HAVCR2 ANTIBODY

SKU: EB07113



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias

kidney injury molecule-3|T cell immunoglobulin mucin-3|T cell immunoglobulin mucin 3|Tim-3|TIMD3|KIM-3|FLJ14428|hepatitis A virus cellular receptor 2|TIM3|HAVCR2

Usage **Summary**

Names

Immunofluorescence: Strong expression of the protein seen in the nucleus and plasma membrane of HepG2 cells. Recommended concentration: 10µg/ml. Flow Cytometry: Flow

cytometric analysis of HepG2 cells. Recommended concentration: 10ug/ml.

Accession

NP_116171.3 ID

Blocking Peptide

EBP07113

Immunogen

Peptide with sequence C-KWYSHSKEKIQN, from the internal region of the protein sequence according to NP 116171.3.





Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

Peptide

C-KWYSHSKEKIQN

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Species

Human, Dog

Reactive

Human Species

Human

84868 **Gene ID**

Product

Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_plus_medium.png

ELISA

Detection

Antibody detection limit dilution 1:64000.

Limit

Western Blot

Approx 35kDa band observed in lysates of cell lines A549 and HepG2 and in preliminary testing of Jurkat and U937 cell lysate (calculated MW of 33.4kDa according to NP_116171.3). Recommended concentration:0.1-1µg/ml. Primary

incubation 1 hour at room temperature.

Application

Type

Pep-ELISA, WB, IF, FC

GALLERY IMAGES



Telephone: (650) 697-3600



