
Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-TNFSF11 / OPGL ANTIBODY





SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

receptor activator of nuclear factor kappa B ligand tumor necrosis factor ligand superfamily, member

11|osteoprotegerin ligand|osteoclast differentiation factor|TNF-related activation-induced **Alias**

Names

cytokine|sOdf|hRANKL2|TRANCE|RANKL|OPGL|ODF|tumor necrosis factor (ligand) superfamily, member

11|TNFSF11

Usage Summary Immunofluorescence: Strong expression of the protein seen in the membranes of MCF7 and NIH3T3 cells. Recommended concentration: 10μg/ml. Flow Cytometry: Flow

cytometric analysis of MCF7 cells. Recommended concentration: 10ug/ml.

Accession

NP_003692.1; NP_143026.1

Blocking Peptide

ID

EBP07163

Immunogen

Peptide with sequence C-DLAKRSKLEAQP, from the internal region of the protein sequence according to NP_003692.1; NP_143026.1.









Product Comments

This antibody is expected to recognise both reported isoforms (NP_003692.1 and NP_143026.1).

Peptide

Sequence

C-DLAKRSKLEAQP

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method

using the immunizing peptide.

Shipping Instructions

Refrigerated

Human

8600

Predicted

Species

Reactive

Human, Mouse

Species

Human

Gene ID

Product

Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png

ELISA

Detection

Antibody detection limit dilution 1:16000.

Limit

Application

Type

Pep-ELISA, IF, FC

GALLERY IMAGES

