








## **GOAT ANTI-TNFSF13 / APRIL ANTIBODY**

**SKU:** EB09379



## **SPECIFICATIONS**

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Unit Size** 

100 μg

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias

tumor necrosis factor-related death ligand-1|tumor necrosis factor ligand superfamily, member 13|tumor necrosis factor ligand superfamily member 13 epsilon|tumor necrosis factor (ligand) superfamily member 13 transcript variant delta|a proliferation inducing ligand|TNF- and APOL-related leukocyte expressed ligand 2|OTTHUMP00000174780|ligand|UNQ383/PRO715|TRDL-1|TALL2|CD256|APRIL|tumor necrosis factor (ligand)

superfamily, member 13|TNFSF13

Accession

**Names** 

NP\_003799.1; NP\_742084.1; NP\_742085.1

**Blocking Peptide** 

ID

EBP09379

**Immunogen** 

Peptide with sequence C-PRARAKLNLSPHGT, from the C-Terminus (near) of the protein sequence according to NP\_003799.1; NP\_742084.1; NP\_742085.1.

**Product** 

This antibody is expected to recognize all isoforms (alpha, beta, gamma: NP\_003799.1; NP\_742084.1;

Comments

NP\_742085.1). This antibody also cross-reacts with human, mouse and dog TNFSF12-TNFSF13 protein (GenelD: 407977, 619441, 479484; 36.6kDa, 45.9kDa, ca. 25.9kDa).





Email: <a href="mailto:customerservice@vectorlabs.com">customerservice@vectorlabs.com</a>

Telephone: (650) 697-3600

**Peptide** C-PRARAKLNLSPHGT

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

**Shipping** 

Instructions

Refrigerated

**Predicted Species** 

Human, Mouse, Rat, Dog, Cow

Reactive

Human **Species** 

Human

8741, 407977

Gene ID Mouse

69583 Gene ID

**Product** 

Rat Gene ID 287437

Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite\_medium.png

**IHC Results** 

In paraffin embedded Human Kidney shows heavy cytoplasm staining in the epithelial cells of DCT.

Recommended concentration, 5-15µg/ml.

**ELISA** 

Detection

Antibody detection limit dilution 1:8000.

Limit

Application

Type

Pep-ELISA, IHC

## GALLERY IMAGES

