



Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-TROP2 ANTIBODY

SKU: EB08839



Telephone: (650) 697-3600



250kDa 150kDa 100kDa

75kDa

50kDa


37kDa

25kDa

20kDa

15kDa







SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym /

membrane component, chromosome 1, surface marker 1 (40kD glycoprotein, identified by monoclonal antibody GA733)|epithelial glycoprotein-1|M1S1|GA733-1|GA733|EGP-1|tumor-associated calcium signal

transducer 2|TACSTD2|TROP2 Names

Accession

NP_002344.2

Blocking

Alias

ID

EBP08839 **Peptide**

Immunogen

Peptide with sequence C-HYEQPTIQIELRQN, from the internal region of the protein sequence according to

NP 002344.2.

Peptide

Sequence

C-HYEOPTIQIELRON

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Shipping

Refrigerated

Instructions

Predicted Species

Human, Mouse, Rat, Dog, Pig, Cow

Reactive

Species

Human

Human

4070 Gene ID

Mouse

56753

Gene ID

Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite medium.png Grade

IHC Results Paraffin embedded Human Tonsil. Recommended concentration: 2.5µg/ml.

ELISA

Detection Antibody detection limit dilution 1:4000.

Limit


Western Approx. 37kDa band observed in Human Prostate lysates (calculated MW of 35.7kDa according to

NP_002344.2). Recommended concentration: 0.2-0.5µg/ml. Primary incubation was 1 hour. Blot

Application

Pep-ELISA, WB, IHC **Type**







GALLERY IMAGES

250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

