


GOAT ANTI-WHIP / WRNIP1 ANTIBODY

SKU: EB05132



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias bA420G6.2|RP11-420G6.2|Werner helicase interacting protein 1|putative helicase RUVBL|FLJ22526|Werner helicase interacting protein|WHIP

Names
Accession ID NP_064520.2; NP_569079.1

Blocking Peptide EBP05132

Immunogen Peptide with sequence EELRGVDFFKQRRC, from the C Terminus of the protein sequence according to NP_064520.2; NP_569079.1.

Product Comments This antibody is expected to recognise isoform 1 (NP_064520.2) and isoform 2 (NP_569079.1).

Peptide Sequence EELRGVDFFKQRRC

Purification Method Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

| | |
|------------------------------|--|
| Shipping Instructions | Refrigerated |
| Predicted Species | Human, Mouse, Rat, Dog, Cow |
| Reactive Species | Human |
| Human Gene ID | 56897 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| IHC Results | In paraffin embedded Human Kidney shows nuclear staining, stronger in DCT compared to PCT. Recommended concentration, 4-6µg/ml. |
| ELISA Detection Limit | Antibody detection limit dilution 1:32000. |
| Western Blot | Approx 90kDa band observed in lysates of cell line Jurkat (calculated MW of 72.1kDa according to NP_064520.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Bish RA, Myers MP., J Biol Chem. 2007 Aug 10;282(32):23184-93; PMID: 17550899). Recommended concentration: 1-3µg/ml. An additional band of unknown identity was also consistently observed at 50kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? |
| Application Type | Pep-ELISA, WB, IHC |

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

