

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06443 - Goat Anti-58KGolgi protein(Internal)/FTCD Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: FTCD, LCHC1, formiminotransferase cyclodeaminase,

formimidoyltransferase cyclodeaminase, LC1 autoantigen

Official Symbol: FTCD

Accession Number(s): NP\_006648.1; NP\_996848.1; NP\_001307341.1

Human GeneID(s): 10841

Important Comments: Variants (NP\_006648.1; NP\_996848.1) encode the same protein.

### **Immunogen**

Peptide with sequence CLREQGRGKDQPGRL, from the internal region of the protein sequence according to NP\_006648.1; NP\_996848.1; NP\_001307341.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 55-58kDa band observed in Human, Mouse and Rat Liver lysates (calculated MW of 58.9kDa according to Human NP\_006648.1, Mouse NP\_543121.1 and Rat NP\_446019.1). Recommended concentration: 0.5-1µg/ml. Primary incubation 1 hour at room temperature.


IHC: Paraffin embedded Human Liver. Recommended concentration: 5-6µg/ml.

**Immunofluorescence:** Expression of the protein seen in the membranes and cytoplasm of Caco-2 cells. Recommended concentration: 10μg/ml. **Flow Cytometry:** A customer reported weak staining on Caco-2 cells at a concentration of 10ug/ml.


#### **Species Reactivity**

Tested: Human, Mouse, Rat


Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow




EB06443 (0.75μg/ml) staining of Human (A) and Mouse (B) and (0.5ug/ml) of Rat © Liver lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB06443 Immunofluorescence analysis of paraformaldehyde fixed Caco-2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06443 (6μg/ml) staining of paraffin embedded Human Liver. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB06443 Negative Control showing staining of paraffin embedded Human Liver, with no primary antibody.