

#### **International Office**

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560

**United States** 

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09741-B - Goat Anti-ACAT1 (aa253-266), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

Principal Names: ACAT1, acetyl-CoA acetyltransferase 1, ACAT, MAT, T2, THIL, acetoacetyl Coenzyme A thiolase, acetoacetyl-CoA thiolase, acetyl-Coenzyme A acetyltransferase 1, mitochondrial acetoacetyl-CoA thiolase Official Symbol: ACAT1 Accession Number(s): NP\_000010.1 Human GeneID(s): <u>38</u> Non-Human GeneID(s): 110446 (mouse), 25014 (rat)

#### Immunogen

Peptide with sequence C-DEEYKRVDFSKVPK., from the internal region of the protein sequence according to NP\_000010.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx 45kDa band observed in Mouse and Rat Liver lysates (calculated MW of 44.8kDa according to Mouse NP\_659033.1 and 44.7kDa according to Rat NP\_058771.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

### **Species Reactivity**

Tested: Human, Mouse, Rat Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



Biotinylated EB09741 (0.1µg/ml) staining of Mouse (A) and Rat (B) Liver lysates (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.