

## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09740 - Goat Anti-ACAT1 (aa257-269) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ACAT1, acetyl-Coenzyme A acetyltransferase 1, ACAT, MAT, T2, THIL, acetoacetyl Coenzyme A thiolase, acetyl-CoA acetyltransferase 1, mitochondrial acetoacetyl-CoA thiolase

**Official Symbol:** ACAT1

**Accession Number(s):** NP\_000010.1

**Human GeneID(s):** [38](#)

**Non-Human GeneID(s):** 110446 (mouse), 25014 (rat)

### Immunogen

Peptide with sequence C-KRVDFSKVPLKKT, from the internal region of the protein sequence according to NP\_000010.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 45kDa band observed in Human Liver lysates and in rodent Kidney and Liver lysates (calculated MW of 45.2kDa according to NP\_000010.1). Recommended concentration: 0.01-0.03µg/ml.

**Immunofluorescence:** Customer finds particulate cytoplasm staining in HeLa


### Species Reactivity

**Tested:** Human, Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Cow



EB09740 (0.01 $\mu$ g/ml) staining of Human Liver lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB09740 (0.01 $\mu$ g/ml) staining of Mouse (A) and Rat (B) kidney lysates, and Mouse (C) and Rat (D) Liver lysates (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.